

Nutrition Management in Neonates: Prolacta Weaning Chart

Prolacta Weaning Protocol Initiated at 33 0/7 Weeks CGA								
Days	Feeding Sequence every 3 hours							
	1	2	3	4	5	6	7	8
Day 1	Prolacta	Prolacta	<i>Sim HMF</i>	Prolacta	Prolacta	<i>Sim HMF</i>	Prolacta	Prolacta
Day 2	Prolacta	<i>Sim HMF</i>	Prolacta	<i>Sim HMF</i>	Prolacta	<i>Sim HMF</i>	Prolacta	<i>Sim HMF</i>
Day 3	<i>Sim HMF</i>	<i>Sim HMF</i>	Prolacta	<i>Sim HMF</i>	<i>Sim HMF</i>	Prolacta	<i>Sim HMF</i>	<i>Sim HMF</i>
Day 4	<i>Sim HMF</i>	<i>Sim HMF</i>	<i>Sim HMF</i>	<i>Sim HMF</i>	<i>Sim HMF</i>	<i>Sim HMF</i>	<i>Sim HMF</i>	<i>Sim HMF</i>

Weaning protocol of Prolacta to Bovine fortifier* at 33 0/7 weeks GA: Add 2 feeding Similac HMF

*For infants born severely SGA who are still <1250 grams at 33 0/7 weeks GA, delay weaning of Prolacta until they reach >1250 grams.

Reviewers:

Created by	Department	Creation Date	Version Date
AAH NICU Standardization Committee	Pediatrics NICU	June 2020	Jan 2024